



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## *On the Expansion of $\phi(x+h)$ .*

By A. W. WHITCOM, M. D.

*Sheboygan Falls, Wisconsin.*

---

THE object of this paper is the development of Taylor's formula, the development of the form of the functional coefficient of  $h$  (usually known as  $\theta$ ), in the remainder in that formula, and the development of the forms of  $\theta_1, \theta_2, \dots$  as they appear in the equations  $(a)', (b)', \dots$

Suppose  $\phi x$  and  $\phi'x$  to be finite and continuous functions for all values of  $x$  from  $x = x'$  to  $x = x' + mh$ ,  $m$  being always positive, and either a constant, or a function of  $x$  and  $h$  of such form as to reduce to a constant when  $h = 0$ . Then will

$$\frac{\phi(x' + mh) - \phi x'}{mh} = \phi'(x' + \theta'h),$$

where  $\theta'$  is between 0 and  $m$ .

Let  $A$  and  $B$  be the algebraically greatest and least values of  $\phi'x$  for values of  $x$  between  $x'$  and  $x' + mh$ .

Put

$$y = Ax - \phi x, \tag{1}$$

and

$$z = \phi x - Bx. \tag{2}$$

Then

$$\frac{dy}{dx} = A - \phi'x,$$

and

$$\frac{dz}{dx} = \phi'x - B,$$

both of which are positive for all values of  $x$  between  $x'$  and  $x' + mh$ , and consequently  $y$  and  $z$  are increasing functions for all values of  $x$  between  $x'$  and  $x' + mh$ .

Let  $x$  take each of the two values  $x'$  and  $x' + mh$  in both (1) and (2). Then

$$y'' = A(x' + mh) - \phi(x' + mh),$$

$$z'' = \phi(x' + mh) - B(x' + mh),$$

$$y' = Ax' - \phi x',$$

and

$$z' = \phi x - Bx'.$$

Then

$$\frac{y'' - y'}{mh} = A - \frac{\phi(x' + mh) - \phi x'}{mh}, \quad \text{and} \quad \frac{z'' - z'}{mh} = \frac{\phi(x' + mh) - \phi x'}{mh} - B.$$

Because  $y$  and  $z$  are increasing functions, and  $m$  is positive,  $y'' - y'$  and  $z'' - z'$  are of the same sign as  $h$ , and the members of the last two equations are positive. Consequently  $\frac{\phi(x' + mh) - \phi x'}{mh}$  is less than  $A$  and greater than  $B$ , and therefore, by reason of the continuity of  $\phi'x$  between  $A$  and  $B$ , is equal to some value of  $\phi'x$  between  $A$  and  $B$  in which  $x$  has some value between  $x'$  and  $x' + mh$ . Let  $x' + \theta'h$  represent this value of  $x$ ,  $\theta'$  being between 0 and  $m$ . Then

$$\frac{\phi(x' + mh) - \phi x'}{mh} = \phi'(x' + \theta'h),$$

or

$$\phi(x + h) = \phi x + mh\phi'(x + \theta'h), \quad (3)$$

which will hold true for all values of  $x$  and  $mh$  within the limits of those giving finite and continuous values in  $\phi x$  and  $\phi'x$ .

Differentiating (3) twice, regarding  $x$  as constant, and in the result putting  $h = 0$ ,

$$(m^2)_{h=0}\phi''x = 2(m\theta')_{h=0}\phi''x.$$

$$\therefore (\theta')_{h=0} = \frac{1}{2}(m)_{h=0}, \quad \text{and} \quad \theta' = \frac{1}{2}(m)_{h=0} + \theta^0,$$

where  $\theta^0 = 0$  when  $h = 0$ . These results obtain whether  $m$  be constant or variable.

Since  $\theta'$  is always between 0 and  $m$ , and  $m$  is always positive, and since  $\theta'$  reduces to the constant  $\frac{1}{2}(m)_{h=0}$  when  $h = 0$ ,  $\theta'$ , like  $m$ , is always positive, and either a constant, or reduces to one when  $h = 0$ . Hence, if  $\phi''x, \phi'''x, \dots, \phi^{2n}x$  are finite and continuous functions the same as  $\phi x$  and  $\phi'x$ , we can form the

following equations, in which  $\theta''$ ,  $\theta''' \dots$  are, like  $m$  and  $\theta'$ , always positive, and either constants, or reduced to constants when  $h = 0$ .

$$\phi(x + mh) = \phi x + mh\phi'(x + \theta'h), \quad (a)$$

where  $\theta'$  is between 0 and  $m$ ;

$$\phi'(x + \theta'h) = \phi'x + \theta'h\phi''(x + \theta''h), \quad (b)$$

where  $\theta''$  is between 0 and  $\theta'$ ;

$$\phi''(x + \theta''h) = \phi''x + \theta''h\phi'''(x + \theta'''h), \quad (c)$$

where  $\theta'''$  is between 0 and  $\theta''$ ;

$$\begin{array}{ccccccccccc} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \phi^{2n-1}(x + \theta^{2n-1}h) = \phi^{2n-1}x + \theta^{2n-1}h\phi^{2n}(x + \theta^{2n}h), & (d) \end{array}$$

where  $\theta^{2n}$  is between 0 and  $\theta^{2n-1}$ .

The same law that makes

$$(\theta')_{h=0} = \frac{1}{2} (m)_{h=0},$$

also makes

$$(\theta'')_{h=0} = \frac{1}{2} (\theta')_{h=0} = \frac{1}{2^2} (m)_{h=0},$$

$$(\theta''')_{h=0} = \frac{1}{2} (\theta'')_{h=0} = \frac{1}{2^3} (m)_{h=0},$$

and in general,

$$(\theta^{2n})_{h=0} = \frac{1}{2^{2n}} (m)_{h=0}.$$

Let  $m = 1$ , and denote what  $\theta'$ ,  $\theta''$ ,  $\theta''' \dots$  become by  $\theta_1$ ,  $\theta_2$ ,  $\theta_3 \dots$ . Then,

$$\phi(x + h) = \phi x + h\phi'(x + \theta_1h), \quad (a)$$

where  $\theta_1$  is between 0 and 1;

$$\phi'(x + \theta_1h) = \phi'x + \theta_1h\phi''(x + \theta_2h), \quad (b)$$

where  $\theta_2$  is between 0 and  $\theta_1$ ;

$$\phi''(x + \theta_2h) = \phi''x + \theta_2h\phi'''(x + \theta_3h), \quad (c)$$

where  $\theta_3$  is between 0 and  $\theta_2$ ;

$$\begin{array}{ccccccccccc} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \phi^{2n-1}(x + \theta_{2n-1}h) = \phi^{2n-1}x + \theta_{2n-1}h\phi^{2n}(x + \theta_{2n}h), & (d) \end{array}$$

where  $\theta_{2n}$  is between 0 and  $\theta_{2n-1}$ .

Further, if  $h = 0$ , then

$$(\theta_1)_{h=0} = \frac{1}{2}, \quad (\theta_2)_{h=0} = \frac{1}{4},$$

$$(\theta_3)_{h=0} = \frac{1}{8} \dots (\theta_{2n})_{h=0} = \frac{1}{2^{2n}}.$$

Forming the successive diff. co. of (3), regarding  $x$  as constant, and in the results putting  $h = 0$ ,

$$(m)_{h=0} \phi'x = (m)_{h=0} \phi'x. \quad (4)$$

$$(m^2)_{h=0} \phi''x = 2(m\theta')_{h=0} \phi''x. \quad (5)$$

$$6 \left( m \frac{dm}{dh} \right)_{h=0} \phi''x + (m^3)_{h=0} \phi'''x = 6 \left( \theta' \frac{dm}{dh} \right)_{h=0} \phi''x + 6 \left( m \frac{d\theta'}{dh} \right)_{h=0} \phi''x + 3(m\theta^2)_{h=0} \phi'''x. \quad (6)$$

$$\begin{aligned} 12 \left( m \frac{d^2m}{dh^2} \right)_{h=0} \phi''x + 12 \left( \frac{dm}{dh} \right)_{h=0}^2 \phi''x + 12 \left( m^2 \frac{dm}{dh} \right)_{h=0} \phi'''x + (m^4)_{h=0} \phi^{IV}x &= 12 \left( \theta' \frac{d^2m}{dh^2} \right)_{h=0} \phi''x \\ &+ 24 \left( \frac{dm}{dh} \frac{d\theta'}{dh} \right)_{h=0} \phi''x + 12 \left( \theta'^2 \frac{dm}{dh} \right)_{h=0} \phi'''x + 12 \left( m \frac{d^2\theta'}{dh^2} \right)_{h=0} \phi''x + 24 \left( m\theta' \frac{d\theta'}{dh} \right)_{h=0} \phi'''x \\ &+ 4(m\theta^3)_{h=0} \phi^{IV}x. \end{aligned} \quad (7)$$

$$\begin{aligned} 20 \left( m \frac{d^3m}{dh^3} \right)_{h=0} \phi''x + 60 \left( \frac{dm}{dh} \frac{d^2m}{dh^2} \right)_{h=0} \phi''x + 30 \left( m^2 \frac{d^2m}{dh^2} \right)_{h=0} \phi'''x + 60 \left[ m \left( \frac{dm}{dh} \right)^2 \right]_{h=0} \phi'''x \\ + 20 \left( m^3 \frac{dm}{dh} \right)_{h=0} \phi^{IV}x + (m^5)_{h=0} \phi^Vx &= 20 \left( \theta' \frac{d^3m}{dh^3} \right)_{h=0} \phi''x + 60 \left( \frac{d^2m}{dh^2} \frac{d\theta'}{dh} \right)_{h=0} \phi''x \\ &+ 30 \left( \theta'^2 \frac{d^2m}{dh^2} \right)_{h=0} \phi'''x + 60 \left[ m \left( \frac{d\theta'}{dh} \right)^2 \right]_{h=0} \phi'''x + 20 \left( \theta'^3 \frac{dm}{dh} \right)_{h=0} \phi^{IV}x + 5(m\theta^4)_{h=0} \phi^Vx \\ &+ 60 \left( \frac{dm}{dh} \frac{d^2\theta'}{dh^2} \right)_{h=0} \phi''x + 120 \left( \theta' \frac{dm}{dh} \frac{d\theta'}{dh} \right)_{h=0} \phi'''x + 20 \left( m \frac{d^3\theta'}{dh^3} \right)_{h=0} \phi''x + 60 \left( m\theta' \frac{d^2\theta'}{dh^2} \right)_{h=0} \phi'''x \\ &+ 60 \left( m\theta'^2 \frac{d\theta'}{dh} \right)_{h=0} \phi^{IV}x. \end{aligned} \quad (8)$$

$$\begin{aligned} 30 \left( m \frac{d^4m}{dh^4} \right)_{h=0} \phi''x + 120 \left( \frac{dm}{dh} \frac{d^3m}{dh^3} \right)_{h=0} \phi''x + 60 \left( m^2 \frac{d^3m}{dh^3} \right)_{h=0} \phi'''x + 90 \left( \frac{d^2m}{dh^2} \right)_{h=0}^2 \phi'''x \\ + 360 \left( m \frac{dm}{dh} \frac{d^2m}{dh^2} \right)_{h=0} \phi'''x + 60 \left( m^2 \frac{d^2m}{dh^2} \right)_{h=0} \phi^{IV}x + 120 \left( \frac{dm}{dh} \right)_{h=0}^3 \phi^{IV}x \\ + 180 \left[ m^2 \left( \frac{dm}{dh} \right)^2 \right]_{h=0} \phi^{IV}x + 30 \left( m^4 \frac{dm}{dh} \right)_{h=0} \phi^Vx + (m^6)_{h=0} \phi^{VI}x &= 30 \left( \theta' \frac{d^4m}{dh^4} \right)_{h=0} \phi''x \\ &+ 120 \left( \frac{d^3m}{dh^3} \frac{d\theta'}{dh} \right)_{h=0} \phi''x + 60 \left( \theta'^2 \frac{d^3m}{dh^3} \right)_{h=0} \phi'''x + 180 \left( \frac{d^2m}{dh^2} \frac{d^2\theta'}{dh^2} \right)_{h=0} \phi''x \end{aligned}$$

$$\begin{aligned}
& + 360 \left( \theta' \frac{d^2 m}{dh^2} \frac{d\theta'}{dh} \right)_{h=0} \phi'''x + 60 \left( \theta'^3 \frac{d^2 m}{dh^2} \right)_{h=0} \phi^{IV}x + 120 \left( \frac{dm}{dh} \frac{d^3 \theta'}{dh^3} \right)_{h=0} \phi''x \\
& + 360 \left( \theta' \frac{dm}{dh} \frac{d^2 \theta'}{dh^2} \right)_{h=0} \phi'''x + 360 \left[ \frac{dm}{dh} \left( \frac{d\theta'}{dh} \right)^2 \right]_{h=0} \phi'''x + 360 \left( \theta'^2 \frac{dm}{dh} \frac{d\theta'}{dh} \right)_{h=0} \phi^{IV}x \\
& + 30 \left( m \frac{d^4 \theta'}{dh^4} \right)_{h=0} \phi''x + 120 \left( m\theta' \frac{d^3 \theta'}{dh^3} \right)_{h=0} \phi'''x + 360 \left( m \frac{d\theta'}{dh} \frac{d^2 \theta'}{dh^2} \right)_{h=0} \phi'''x \\
& + 180 \left( m\theta'^2 \frac{d^2 \theta'}{dh^2} \right)_{h=0} \phi^{IV}x + 30 \left( \theta'^4 \frac{dm}{dh} \right)_{h=0} \phi^Vx + 360 \left[ m\theta' \left( \frac{d\theta'}{dh} \right)^2 \right]_{h=0} \phi^{IV}x \\
& + 120 \left( m\theta'^3 \frac{d\theta'}{dh} \right)_{h=0} \phi^Vx + 6 (m\theta'^5)_{h=0} \phi^{VI}x.
\end{aligned} \tag{9}$$

If in (4), (5), (6), (7), (8), and (9) we put  $m = 1$ , and write  $\theta_1$  for  $\theta'$ , we obtain the same results as when  $h = 0$  in the successive diff. co., with respect to  $h$ , of  $(a)'$ . Hence,

$$\phi'x = \phi'x. \tag{10}$$

$$\phi''x = 2(\theta_1)_{h=0} \phi''x. \tag{11}$$

$$\phi'''x = 6 \left( \frac{d\theta_1}{dh} \right)_{h=0} \phi''x + 3(\theta_1^2)_{h=0} \phi'''x. \tag{12}$$

$$\phi^{IV}x = 12 \left( \frac{d^2 \theta_1}{dh^2} \right)_{h=0} \phi''x + 24 \left( \theta_1 \frac{d\theta_1}{dh} \right)_{h=0} \phi'''x + 4(\theta_1^3)_{h=0} \phi^{IV}x. \tag{13}$$

$$\begin{aligned}
\phi^Vx &= 20 \left( \frac{d^3 \theta_1}{dh^3} \right)_{h=0} \phi''x + 60 \left( \theta_1 \frac{d^2 \theta_1}{dh^2} \right)_{h=0} \phi'''x + 60 \left( \frac{d\theta_1}{dh} \right)_{h=0}^2 \phi'''x + 60 \left( \theta_1^2 \frac{d\theta_1}{dh} \right)_{h=0} \phi^{IV}x \\
&+ 5(\theta_1^4)_{h=0} \phi^Vx.
\end{aligned} \tag{14}$$

$$\begin{aligned}
\phi^{VI}x &= 30 \left( \frac{d^4 \theta_1}{dh^4} \right)_{h=0} \phi''x + 120 \left( \theta_1 \frac{d^3 \theta_1}{dh^3} \right)_{h=0} \phi'''x + 360 \left( \frac{d\theta_1}{dh} \frac{d^2 \theta_1}{dh^2} \right)_{h=0} \phi'''x + 180 \left( \theta_1^2 \frac{d^2 \theta_1}{dh^2} \right)_{h=0} \phi^{IV}x \\
&+ 360 \left[ \theta_1 \left( \frac{d\theta_1}{dh} \right)^2 \right]_{h=0} \phi^{IV}x + 120 \left( \theta_1^3 \frac{d\theta_1}{dh} \right)_{h=0} \phi^Vx + 6(\theta_1^5)_{h=0} \phi^{VI}x.
\end{aligned} \tag{15}$$

Similarly,

$$\begin{aligned}
\phi^{VII}x &= 42 \left( \frac{d^5 \theta_1}{dh^5} \right)_{h=0} \phi''x + 210 \left( \theta_1 \frac{d^4 \theta_1}{dh^4} \right)_{h=0} \phi'''x + 840 \left( \frac{d\theta_1}{dh} \frac{d^3 \theta_1}{dh^3} \right)_{h=0} \phi'''x + 420 \left( \theta_1^2 \frac{d^3 \theta_1}{dh^3} \right)_{h=0} \phi^{IV}x \\
&+ 630 \left( \frac{d^2 \theta_1}{dh^2} \right)_{h=0}^2 \phi''x + 2520 \left( \theta_1 \frac{d\theta_1}{dh} \frac{d^2 \theta_1}{dh^2} \right)_{h=0} \phi^{IV}x + 840 \left( \frac{d\theta_1}{dh} \right)_{h=0}^3 \phi^{IV}x \\
&+ 1260 \left[ \theta_1^2 \left( \frac{d\theta_1}{dh} \right)^2 \right]_{h=0} \phi^Vx + 420 \left( \theta_1^3 \frac{d^2 \theta_1}{dh^2} \right)_{h=0} \phi^Vx + 210 \left( \theta_1^4 \frac{d\theta_1}{dh} \right)_{h=0} \phi^{VI}x \\
&+ 7(\theta_1^6)_{h=0} \phi^{VII}x.
\end{aligned} \tag{16}$$

Equation  $(a)'$  reduces to  $\phi x = \phi x$ , when  $h = 0$ , but its diff. co., with respect to  $h$ , reduces to  $\phi'x = \phi'x$ , by (10). Hence by one differentiation with respect to  $h$ ,  $h$  has been eliminated from one term in  $h\phi'(x + \theta_1 h)$ . Consequently one term in  $h\phi'(x + \theta_1 h)$  is  $h\phi'x$ .

Again: when  $h = 0$ , the second diff. co. of  $(a)'$  with respect to  $h$ , reduces to  $\phi''x = 2(\theta_1)_{h=0}\phi''x$ , by (11). Consequently by two differentiations with respect to  $h$ ,  $\frac{h^2}{2}$  has been eliminated from a second term in  $h\phi'(x + \theta_1 h)$ , and a second term in that expression is

$$\frac{h^2}{2} \phi''x = \frac{h^2}{2} [2(\theta_1)_{h=0}\phi''x].$$

In like manner by three differentiations with respect to  $h$ , and in the result putting  $h = 0$ , (12) is obtained from  $(a)'$ , where it is plain  $\frac{h^3}{2.3}$  has been eliminated from a third term in  $h\phi'(x + \theta_1 h)$ , and a third term in that expression is

$$\frac{h^3}{2.3} \phi'''x = \frac{h^3}{2.3} \left[ 6 \left( \frac{d\theta_1}{dh} \right)_{h=0} \phi''x + 3(\theta_1^2)_{h=0} \phi'''x \right].$$

Additional terms may be found in the same way from (13), (14) . . . ., and with the foregoing ones substituted for  $h\phi'(x + \theta_1 h)$  in  $(a)'$ , giving

$$\phi(x+h) = \phi x + h\phi'x + \frac{h^2}{2} \phi''x + \frac{h^3}{2.3} \phi'''x + \dots \frac{h^n}{[n]} \phi^n x + \dots \quad (17)$$

If, in (17),  $\frac{h^n}{[n]} \phi^n(x + \theta h)$  denote the sum of all the terms in the right member after the first  $n$ , and again  $\frac{h^m}{[m]} R$  denote the sum of all the terms after the first  $m$ ,  $m$  being any integer between  $n$  and  $2n$ , we can write the two equations

$$\phi(x+h) = \phi x + h\phi'x + \frac{h^2}{2} \phi''x + \frac{h^3}{2.3} \phi'''x + \dots \frac{h^{n-1}}{[n-1]} \phi^{n-1}x + \frac{h^n}{[n]} \phi^n(x + \theta h). \quad (18)$$

$$\begin{aligned} \phi(x+h) &= \phi x + h\phi'x + \frac{h^2}{2} \phi''x + \frac{h^3}{2.3} \phi'''x + \dots \frac{h^n}{[n]} \phi^n x + \frac{h^{n+1}}{[n+1]} \phi^{n+1}x \\ &+ \dots \frac{h^{m-1}}{[m-1]} \phi^{m-1}x + \frac{h^m}{[m]} R. \end{aligned} \quad (19)$$

From (18) and (19),

$$\phi^n(x + \theta h) = \phi^n x + \frac{h}{(n+1)} \phi^{n+1}x + \frac{h^2}{(n+1)(n+2)} \phi^{n+2}x + \dots \frac{h^{m-n}}{(n+1)(n+2)\dots m} R. \quad (20)$$

By differentiating (18)  $n$  times, regarding  $x$  as constant,

$$\begin{aligned} \phi^n(x+h) &= \phi^n(x + \theta h) + nh \frac{d\phi^n(x + \theta h)}{dh} + \frac{n(n-1)h^2}{2} \frac{d^2\phi^n(x + \theta h)}{dh^2} \\ &+ \dots \frac{h^n}{[n]} \frac{d^n\phi^n(x + \theta h)}{dh^n}. \end{aligned} \quad (21)$$

From (18), when  $n=0$ ,

$$\phi(x+h) = \frac{h^0}{1} \phi^0(x+\theta h) = \phi^0(x+\theta h), \quad \text{and} \quad \theta = 1. \quad (22)$$

If  $n = \infty$ , then, by (20),

$$\phi^n(x+\theta h) = \phi^n x, \quad \text{and} \quad \theta = 0. \quad (23)$$

From (20) and (21) we find the quantity  $\phi^n(x+\theta h)$  can never equal either  $\phi^n(x+h)$ , or  $\phi^n x$ , unless  $n$  has such a value as to cause all the terms after the first, in the right members of (20) and (21), to vanish.

This can only happen in (20) when  $n$  is infinite, and in (21) when  $n$  is zero. Hence for finite integral values of  $n$ ,  $\theta$  is a positive proper fraction, or  $\theta$  is between zero and a unit.

Differentiating (20), regarding  $x$  as constant,

$$\begin{aligned} \theta \phi^{n+1}(x+\theta h) + \frac{d\theta}{dh} h \phi^{n+1}(x+\theta h) &= \frac{1}{n+1} \phi^{n+1} x + \frac{2h}{(n+1)(n+2)} \phi^{n+2} x \\ &+ \frac{3h^2}{(n+1)(n+2)(n+3)} \phi^{n+3} x + \dots \frac{(m-n)h^{m-n-1}}{(n+1)\dots m} R' + \frac{h^{m-n}}{(n+1)(n+2)\dots m} \frac{dR'}{dh}. \end{aligned} \quad (24)$$

In (24), when  $h=0$ ,

$$(\theta)_{h=0} \phi^{n+1} x = \frac{1}{n+1} \phi^{n+1} x, \quad \text{and} \quad (\theta)_{h=0} = \frac{1}{n+1}. \quad (25)$$

Continuing the differentiation of (24), regarding  $x$  as constant, and in the results putting  $h=0$ ,

$$2 \left( \frac{d\theta}{dh} \right)_{h=0} \phi^{n+1} x + (\theta^2)_{h=0} \phi^{n+2} x = \frac{2\phi^{n+2} x}{(n+1)(n+2)}. \quad (26)$$

$$3 \left( \frac{d^2\theta}{dh^2} \right)_{h=0} \phi^{n+1} x + 6 \left( \theta \frac{d\theta}{dh} \right)_{h=0} \phi^{n+2} x + (\theta^3)_{h=0} \phi^{n+3} x = \frac{2.3\phi^{n+3} x}{(n+1)(n+2)(n+3)} \quad (27)$$

$$\begin{aligned} 4 \left( \frac{d^3\theta}{dh^3} \right)_{h=0} \phi^{n+1} x + 12 \left( \theta \frac{d^2\theta}{dh^2} \right)_{h=0} \phi^{n+2} x + 12 \left( \frac{d\theta}{dh} \right)_{h=0}^2 \phi^{n+2} x + 12 \left( \theta^2 \frac{d\theta}{dh} \right)_{h=0} \phi^{n+3} x \\ + (\theta^4)_{h=0} \phi^{n+4} x = \frac{2.3.4\phi^{n+4} x}{(n+1)(n+2)\dots(n+4)}. \end{aligned} \quad (28)$$

. . . . .

From (25) and (26),

$$\left( \frac{d\theta}{dh} \right)_{h=0} = \left( \frac{2(n+1) - (n+2)}{2(n+1)^2 \cdot (n+2)} \right) \left( \frac{\phi^{n+2} x}{\phi^{n+1} x} \right). \quad (29)$$

Again, from (27), by the aid of (25) and (29),

$$\left( \frac{d^2\theta}{dh^2} \right)_{h=0} = \left( \frac{2.3(n+1)^2 - (n+2)(n+3)}{3(n+1)^3 \cdot (n+2)(n+3)} \right) \left( \frac{\phi^{n+3} x}{\phi^{n+1} x} \right) - \left( \frac{2(n+1) - (n+2)}{(n+1)^3 \cdot (n+2)} \right) \left( \frac{\phi^{n+2} x}{\phi^{n+1} x} \right)^2. \quad (30)$$



In like manner it may be shown, from (28), that

$$\begin{aligned} \left(\frac{d^3\theta}{dh^3}\right) &= \left(\frac{2 \cdot 3 \cdot 4(n+1)^3 - (n+2)(n+3)(n+4)}{2^2(n+1)^4 \cdot (n+2)(n+3)(n+4)}\right) \left(\frac{\phi^{n+4}x}{\phi^{n+1}x}\right) - \left(\frac{2 \cdot 3(n+1) - 3(n+2)}{2(n+1)^4 \cdot (n+2)}\right) \\ &+ \frac{2 \cdot 3^2(n+1)^2 - 3(n+2)(n+3)}{3(n+1)^4 \cdot (n+2)(n+3)} \left(\frac{\phi^{n+3}x}{\phi^{n+1}x} \cdot \frac{\phi^{n+2}x}{\phi^{n+1}x}\right) - \left(\frac{2^2 \cdot 3(n+1)^2 - 2^2 \cdot 3(n+1)(n+2) + 3(n+2)^2}{2^2(n+1)^4 \cdot (n+2)^2}\right) \\ &- \frac{2 \cdot 3(n+1) - 3(n+2)}{(n+1)^4 \cdot (n+2)} \left(\frac{\phi^{n+2}x}{\phi^{n+1}x}\right)^2. \end{aligned} \quad (31)$$

In the same way, the values of  $\left(\frac{d^4\theta}{dh^4}\right)_{h=0}$ ,  $\left(\frac{d^5\theta}{dh^5}\right)_{h=0}$ , . . . may be found.

From (25),  $\theta = \frac{1}{n+1} + \phi$ , where  $\phi = 0$  when  $h = 0$ ; but by one differentiation with respect to  $h$ , and in the result putting  $h = 0$ ,

$$\left(\frac{d\theta}{dh}\right)_{h=0} = \left(\frac{d\phi}{dh}\right)_{h=0} = \left(\frac{2(n+1) - (n+2)}{2(n+1)^2 \cdot (n+2)}\right) \left(\frac{\phi^{n+2}x}{\phi^{n+1}x}\right),$$

by (29). Consequently, by one differentiation with respect to  $h$ ,  $h$  has been eliminated from one term in  $\phi$ , and also from a second term in  $\theta$ . Hence, before differentiation,

$$\theta = \frac{1}{n+1} + h \left(\frac{d\theta}{dh}\right)_{h=0} + \phi' = \frac{1}{n+1} + h \left[\frac{2(n+1) - (n+2)}{2(n+1)^2 \cdot (n+2)} \left(\frac{\phi^{n+2}x}{\phi^{n+1}x}\right)\right] + \phi',$$

where  $\phi'$  and  $\left(\frac{d\phi'}{dh}\right)$  both vanish when  $h = 0$ . Differentiating the last equation twice, regarding  $x$  as constant, and in the result putting  $h = 0$ ,  $\left(\frac{d^2\theta_1}{dh^2}\right)_{h=0} = \left(\frac{d^2\phi'}{dh^2}\right)_{h=0}$  = the right member (30). Hence by two differentiations with respect to  $h$ ,  $\frac{h^2}{2}$  has been eliminated from a term in  $\phi'$ , and from a third term in  $\theta$ . Consequently,

$$\begin{aligned} \theta &= \frac{1}{n+1} + h \left(\frac{d\theta}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta}{dh^2}\right)_{h=0} + \phi'' = \frac{1}{n+1} + h \left[\left(\frac{2(n+1) - (n+2)}{2(n+1)^2 \cdot (n+2)}\right) \left(\frac{\phi^{n+2}x}{\phi^{n+1}x}\right)\right] \\ &+ \frac{h^2}{2} \left[\left(\frac{2 \cdot 3(n+1)^2 - (n+2)(n+3)}{3(n+1)^3 \cdot (n+2) \cdot (n+3)}\right) \left(\frac{\phi^{n+3}x}{\phi^{n+1}x}\right) - \left(\frac{2(n+1) - (n+2)}{(n+1)^3 \cdot (n+2)}\right) \left(\frac{\phi^{n+2}x}{\phi^{n+1}x}\right)^2\right] + \phi'', \end{aligned}$$

where  $\phi''$ ,  $\frac{d\phi''}{dh}$ , and  $\frac{d^2\phi''}{dh^2}$  vanish when  $h = 0$ . In like manner additional terms may be found, and the general form of  $\theta$  shown to be,

$$\theta = \frac{1}{n+1} + h \left(\frac{d\theta}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta}{dh^2}\right)_{h=0} + \frac{h^3}{2 \cdot 3} \left(\frac{d^3\theta}{dh^3}\right)_{h=0} + \dots + \frac{h^{n-1}}{[n-1]} \left(\frac{d^{n-1}\theta}{dh^{n-1}}\right)_{h=0} + \dots, \quad (32)$$

where  $\left(\frac{d\theta}{dh}\right)_{h=0}$ ,  $\left(\frac{d^2\theta}{dh^2}\right)_{h=0}$ ,  $\dots$  are expressible in terms  $\phi^{n+1}x$ ,  $\phi^{n+2}x$ ,  $\dots$   $\phi^{2n}x$ , as in (29), (30), (31).

From (11), (12),  $\dots$  (16), it may readily be shown that

$$\left(\frac{d\theta_1}{dh}\right)_{h=0} = \frac{1}{24} \frac{\phi'''x}{\phi''x}. \quad (33)$$

$$\left(\frac{d^2\theta_1}{dh^2}\right)_{h=0} = \frac{1}{24} \frac{\phi^{IV}x}{\phi''x} - \frac{1}{24} \left(\frac{\phi'''x}{\phi''x}\right)^2. \quad (34)$$

$$\left(\frac{d^3\theta_1}{dh^3}\right)_{h=0} = \frac{11}{320} \frac{\phi^Vx}{\phi''x} - \frac{3}{32} \frac{\phi^{IV}x}{\phi''x} \frac{\phi'''x}{\phi''x} + \frac{11}{192} \left(\frac{\phi'''x}{\phi''x}\right)^3. \quad (35)$$

$$\left(\frac{d^4\theta_1}{dh^4}\right)_{h=0} = \frac{13}{480} \frac{\phi^{VI}x}{\phi''x} - \frac{43}{480} \frac{\phi^Vx}{\phi''x} \frac{\phi'''x}{\phi''x} + \frac{7}{32} \frac{\phi^{IV}x}{\phi''x} \left(\frac{\phi'''x}{\phi''x}\right)^2 - \frac{1}{16} \left(\frac{\phi^{IV}x}{\phi''x}\right)^2 - \frac{3}{32} \left(\frac{\phi'''x}{\phi''x}\right)^4, \quad (36)$$

$$\begin{aligned} \left(\frac{d^5\theta_1}{dh^5}\right)_{h=0} = & \frac{19}{896} \frac{\phi^{VII}x}{\phi''x} - \frac{31}{384} \frac{\phi^{VI}x}{\phi''x} \frac{\phi'''x}{\phi''x} + \frac{15}{64} \frac{\phi^Vx}{\phi''x} \left(\frac{\phi'''x}{\phi''x}\right)^2 - \frac{55}{108} \frac{\phi^{IV}x}{\phi''x} \left(\frac{\phi'''x}{\phi''x}\right)^3 + \frac{5}{16} \left(\frac{\phi^{IV}x}{\phi''x}\right)^2 \frac{\phi'''x}{\phi''x} \\ & - \frac{53}{384} \frac{\phi^Vx}{\phi''x} \frac{\phi^{IV}x}{\phi''x} + \frac{185}{1152} \left(\frac{\phi'''x}{\phi''x}\right)^5. \end{aligned} \quad (37)$$

The value of  $\theta_1$  may be found from those of

$$\left(\frac{d\theta_1}{dh}\right)_{h=0}, \left(\frac{d^2\theta_1}{dh^2}\right)_{h=0}, \left(\frac{d^3\theta_1}{dh^3}\right)_{h=0}, \dots$$

in the same manner that the value of  $\theta$  was derived from those of

$$\left(\frac{d\theta}{dh}\right)_{h=0}, \left(\frac{d^2\theta}{dh^2}\right)_{h=0}, \dots$$

Hence,

$$\theta_1 = \frac{1}{2} + h \left[ \frac{1}{24} \frac{\phi'''x}{\phi''x} \right] + \frac{h^2}{2} \left[ \frac{1}{24} \frac{\phi^{IV}x}{\phi''x} - \frac{1}{24} \left(\frac{\phi'''x}{\phi''x}\right)^2 \right] + \dots, \quad (38)$$

or,

$$\theta_1 = \frac{1}{2} + h \left(\frac{d\theta_1}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta_1}{dh^2}\right)_{h=0} + \frac{h^3}{2 \cdot 3} \left(\frac{d^3\theta_1}{dh^3}\right)_{h=0} + \dots + \frac{h^n}{n} \left(\frac{d^n\theta_1}{dh^n}\right)_{h=0} + \dots, \quad (39)$$

where

$$\left(\frac{d\theta_1}{dh}\right)_{h=0}, \left(\frac{d^2\theta_1}{dh^2}\right)_{h=0}, \dots, \left(\frac{d^n\theta_1}{dh^n}\right)_{h=0}$$

are expressible in terms of  $\phi''x$ ,  $\phi'''x$ ,  $\dots$   $\phi^{n+2}x$ .

If in (5), (6), (7), (8) and (9) we write  $\theta_1$  for  $m$ ,  $\theta_2$  for  $\theta$ , and advance each order of diff. co. of  $x$  to the next higher, we obtain the same results as when

$h = 0$  in the successive diff. co. with respect to  $h$ , of  $(b)'$ , and these results by easy reductions give

$$\left(\frac{d\theta_2}{dh}\right)_{h=0} = \frac{1}{96} \frac{\phi^{IV}x}{\phi'''x} + \frac{1}{48} \frac{\phi'''x}{\phi''x}. \quad (40)$$

$$\left(\frac{d^2\theta_2}{dh^2}\right)_{h=0} = \frac{1}{192} \frac{\phi^Vx}{\phi'''x} - \frac{1}{192} \left(\frac{\phi^{IV}x}{\phi'''x}\right)^2 + \frac{7}{288} \frac{\phi^{IV}x}{\phi''x} - \frac{1}{48} \left(\frac{\phi'''x}{\phi''x}\right)^2. \quad (41)$$

$$\begin{aligned} \left(\frac{d^3\theta_2}{dh^3}\right)_{h=0} = & \frac{11}{5120} \frac{\phi^{VI}x}{\phi'''x} - \frac{3}{512} \frac{\phi^Vx}{\phi'''x} \frac{\phi^{IV}x}{\phi''x} + \frac{11}{3072} \left(\frac{\phi^{IV}x}{\phi'''x}\right)^2 + \frac{27}{1280} \frac{\phi^Vx}{\phi''x} + \frac{1}{768} \frac{\phi^{IV}x}{\phi''x} \frac{\phi^{IV}x}{\phi'''x} \\ & - \frac{119}{2304} \frac{\phi^{IV}x}{\phi''x} \frac{\phi'''x}{\phi''x} + \frac{11}{384} \left(\frac{\phi'''x}{\phi''x}\right)^3. \end{aligned} \quad (42)$$

From these values of  $\left(\frac{d\theta_2}{dh}\right)_{h=0}$ ,  $\left(\frac{d^2\theta_2}{dh^2}\right)_{h=0}$ ,  $\dots$ , we find

$$\begin{aligned} \theta_2 = & \frac{1}{4} + h \left[ \frac{1}{96} \frac{\phi^{IV}x}{\phi'''x} + \frac{1}{48} \frac{\phi'''x}{\phi''x} \right] + \frac{h^2}{2} \left[ \frac{1}{192} \frac{\phi^Vx}{\phi'''x} - \frac{1}{192} \left(\frac{\phi^{IV}x}{\phi'''x}\right)^2 + \frac{7}{288} \frac{\phi^{IV}x}{\phi''x} - \frac{1}{48} \left(\frac{\phi'''x}{\phi''x}\right)^2 \right] \\ & + \dots \end{aligned} \quad (43)$$

or,

$$\theta_2 = \frac{1}{4} + h \left(\frac{d\theta_2}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta_2}{dh^2}\right)_{h=0} + \frac{h^3}{2 \cdot 3} \left(\frac{d^3\theta_2}{dh^3}\right)_{h=0} + \dots + \frac{h^n}{[n]} \left(\frac{d^n\theta_2}{dh^n}\right)_{h=0} + \dots, \quad (44)$$

where

$$\left(\frac{d\theta_2}{dh}\right)_{h=0}, \left(\frac{d^2\theta_2}{dh^2}\right)_{h=0}, \dots, \left(\frac{d^n\theta_2}{dh^n}\right)_{h=0}$$

are expressible in terms of  $\phi''x$ ,  $\phi'''x$ ,  $\dots$ ,  $\phi^{n+3}x$ , as in (40), (41), (42).

If in (5), (6), (7), (8) and (9) we write  $\theta_2$  for  $m$ ,  $\theta_3$  for  $\theta$ ,  $\phi'''x$  for  $\phi'x$ ,  $\phi^{IV}x$  for  $\phi''x$ , and so on, then reduce the results, we get

$$\left(\frac{d\theta_3}{dh}\right)_{h=0} = \frac{1}{384} \frac{\phi^Vx}{\phi^{IV}x} + \frac{1}{192} \frac{\phi^{IV}x}{\phi'''x} + \frac{1}{96} \frac{\phi'''x}{\phi''x}. \quad (45)$$

$$\begin{aligned} \left(\frac{d^2\theta_3}{dh^2}\right)_{h=0} = & \frac{1}{1536} \frac{\phi^{VI}x}{\phi^{IV}x} - \frac{1}{1536} \left(\frac{\phi^Vx}{\phi^{IV}x}\right)^2 + \frac{7}{2304} \frac{\phi^Vx}{\phi'''x} - \frac{1}{384} \left(\frac{\phi^{IV}x}{\phi'''x}\right)^2 + \frac{7}{576} \frac{\phi^{IV}x}{\phi''x} + \frac{1}{1152} \frac{\phi^Vx}{\phi^{IV}x} \frac{\phi'''x}{\phi''x} \\ & - \frac{1}{96} \left(\frac{\phi'''x}{\phi''x}\right)^2. \end{aligned} \quad (46)$$

From these values of  $\left(\frac{d\theta_3}{dh}\right)_{h=0}$ ,  $\left(\frac{d^2\theta_3}{dh^2}\right)_{h=0}$ ,  $\dots$ , we find

$$\theta_3 = \frac{1}{8} + h \left(\frac{d\theta_3}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta_3}{dh^2}\right)_{h=0} + \dots + \frac{h^n}{[n]} \left(\frac{d^n\theta_3}{dh^n}\right)_{h=0} + \dots \quad (47)$$

where  $\left(\frac{d\theta_3}{dh}\right)_{h=0}$ ,  $\left(\frac{d^2\theta_3}{dh^2}\right)_{h=0}$ ,  $\dots$  are expressible in terms of  $\phi''x$ ,  $\phi'''x$ ,  $\dots$   $\phi^{n+4}x$ , as in (45), (46).

In general, it may be shown that

$$\theta_n = \frac{1}{2^n} + h \left(\frac{d\theta_n}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta_n}{dh^2}\right)_{h=0} + \frac{h^3}{2 \cdot 3} \left(\frac{d^3\theta_n}{dh^3}\right)_{h=0} + \dots + \frac{h^{n-1}}{[n-1]} \left(\frac{d^{n-1}\theta_n}{dh^{n-1}}\right)_{h=0} + \dots, \quad (48)$$

where  $\left(\frac{d\theta_n}{dh}\right)_{h=0}$ ,  $\left(\frac{d^2\theta_n}{dh^2}\right)_{h=0}$ ,  $\dots$ ,  $\left(\frac{d^{n-1}\theta_n}{dh^{n-1}}\right)_{h=0}$  can be expressed in terms of  $\phi''x$ ,  $\phi'''x$ ,  $\dots$   $\phi^{2n}x$ , as in preceding cases.

Eliminating  $\phi'(x + \theta_1 h)$ ,  $\phi''(x + \theta_2 h)$ ,  $\dots$   $\phi^{n-1}(x + \theta_{n-1} h)$  from (a)', (b)',  $\dots$ , there results

$$\phi(x+h) = \phi x + h\phi'x + \theta_1 h^2 \phi''x + \theta_1 \theta_2 h^3 \phi'''x + \dots \theta_1 \theta_2 \theta_3 \dots \theta_{n-1} h^n \phi^n(x + \theta_n h). \quad (49)$$

Since  $\theta_1$  contains  $h$ ,  $h^2$ ,  $h^3$ ,  $\dots$  as factors, and does not contain  $h$  in any other form, and since the same holds true in the expressions giving the values of  $\theta_2$ ,  $\theta_3$ ,  $\dots$ , it is plain that (49) will contain  $h$  as a factor in the regular ascending integral powers,  $h$ ,  $h^2$ ,  $h^3$ ,  $\dots$ , and will not contain  $h$  in any other form, when  $\theta_1$ ,  $\theta_2$ ,  $\dots$  are replaced by their values in terms of  $x$  and  $h$ .

From (39), (44), (47), and (48), we readily find

$$\theta_1 h^2 \phi''x = h^2 \phi''x \left[ \frac{1}{2} + h \left(\frac{d\theta_1}{dh}\right)_{h=0} + \frac{h^2}{2} \left(\frac{d^2\theta_1}{dh^2}\right)_{h=0} + \frac{h^3}{2 \cdot 3} \left(\frac{d^3\theta_1}{dh^3}\right)_{h=0} \right] + \text{terms containing } h^6, h^7, \dots \quad (50)$$

$$\theta_1 \theta_2 h^3 \phi'''x = h^3 \phi'''x \left[ \frac{1}{8} + \frac{h}{4} \left(\frac{d\theta_1}{dh} \frac{2d\theta_2}{dh}\right)_{h=0} + \frac{h^2}{8} \left(\frac{d^2\theta_1}{dh^2} + \frac{8d\theta_1 d\theta_2}{dh dh} + \frac{2d^2\theta_2}{dh^2}\right)_{h=0} \right] + \text{terms containing } h^8, h^7, \dots \quad (51)$$

$$\theta_1 \theta_2 \theta_3 h^4 \phi^{iv}x = h^4 \phi^{iv}x \left[ \frac{1}{16} + \frac{h}{32} \left(\frac{d\theta_1}{dh} + \frac{2d\theta_2}{dh} + \frac{4d\theta_3}{dh}\right)_{h=0} \right] + \text{terms containing } h^6, h^7, \dots \quad (52)$$

$$\theta_1 \theta_2 \theta_3 \theta_4 h^5 \phi^v x = h^5 \phi^v x \left[ \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{1}{8} \cdot \frac{1}{16} \right] + \text{terms containing } h^8, h^7, \dots \quad (53)$$

In (50), (51), (52), (53), replacing the several diff. co. of  $\theta_1$ ,  $\theta_2$ ,  $\dots$  by their values in terms of  $x$  and  $h$ , and substituting the results for  $\theta_1 h^2 \phi''x$ ,  $\theta_1 \theta_2 h^3 \phi'''x$ ,  $\dots$  in (49), we get

$$\begin{aligned}
 \phi(x+h) = & \phi x + h\phi'x + \frac{h^2}{2}\phi''x + \frac{h^3}{24}\phi'''x + \frac{h^4}{48}\phi^{\text{IV}}x - \frac{h^4}{48}\frac{(\phi''x)^2}{\phi''x} + \frac{11h^5}{1920}\phi^{\text{V}}x - \frac{3h^5}{192}\frac{\phi''x}{\phi''x} + \frac{11h^5}{1152}\frac{(\phi''x)^3}{(\phi''x)^2} \\
 & + \frac{h^3}{8}\phi'''x + \frac{h^4}{192}\phi^{\text{IV}}x + \frac{h^4}{96}\frac{(\phi''x)^2}{\phi''x} + \frac{h^5}{768}\phi^{\text{V}}x + \frac{h^5}{192}\frac{\phi''x}{\phi''x} - \frac{h^5}{192}\frac{(\phi''x)^3}{(\phi''x)^2} \\
 & + \frac{h^4}{64}\phi^{\text{IV}}x + \frac{h^4}{96}\frac{(\phi''x)^2}{\phi''x} + \frac{h^5}{3072}\phi^{\text{V}}x + \frac{h^5}{2304}\frac{\phi''x}{\phi''x} + \frac{h^5}{1152}\frac{(\phi''x)^3}{(\phi''x)^2} \\
 & + \frac{h^5}{1024}\phi^{\text{V}}x + \frac{7h^5}{1152}\frac{\phi''x}{\phi''x} - \frac{h^5}{192}\frac{(\phi''x)^3}{(\phi''x)^2} \\
 & + \frac{h^5}{768}\frac{\phi''x}{\phi''x} \\
 & + \frac{h^5}{768}\frac{\phi''x}{\phi''x} \\
 & + \frac{h^5}{768}\frac{\phi''x}{\phi''x} \\
 & - \frac{h^5}{768}\frac{(\phi^{\text{IV}}x)^2}{\phi''x} + \text{terms containing } h^6, h^7, \dots, \\
 & + \frac{h^5}{1536}\frac{(\phi^{\text{IV}}x)^2}{\phi''x} \\
 & + \frac{h^5}{1536}\frac{(\phi^{\text{IV}}x)^2}{\phi''x}.
 \end{aligned}$$

Uniting terms,

$$\begin{aligned}
 \phi(x+h) = & \phi x + h\phi'x + \frac{h^2}{2}\phi''x + \frac{h^3}{2.3}\phi'''x + \frac{h^4}{2.3.4}\phi^{\text{IV}}x + \frac{h^5}{2.3.4.5}\phi^{\text{V}}x + \text{terms} \\
 & \text{containing } h^6, h^7, \dots
 \end{aligned}$$

By extending the work, it may be shown that  $\phi(x+h) = \phi x + h\phi'x + \frac{h^2}{2}\phi''x + \frac{h^3}{2.3}\phi'''x + \dots + \frac{h^n}{[n]}\phi^n x + \text{such terms in } \theta_1 h^2 \phi''x, \theta_1 \theta_2 h^3 \phi'''x, \dots, \theta_1 \theta_2, \dots, \theta_{n-1} h^n \phi^n(x + \theta_n h)$  as contain  $h^{n+1}, h^{n+2}, \dots$ , but no power of  $h$  less than  $h^{n+1}$ . The sum of all the terms in the right member of the last equation, after the first  $n$ , may be denoted by  $\frac{h^n}{[n]}\phi^n(x + \theta h)$ , giving (18), and the value of  $\theta$  found as before.